

## Exploring the role of physical inactivity in gestational diabetes mellitus: a systematic review of evidence

Seveka Bali<sup>1(A,B,E,F)</sup>, Sonia Singh<sup>1(A,D,E,F)</sup>, Aashima Arora<sup>2(B,C,E,F)</sup>

<sup>1</sup>Physiotherapy Department, Punjabi University, Patiala, India

<sup>2</sup>Department of Obstetrics and Gynecology, Postgraduate Institute of Medical Education and Research, Chandigarh, India

### Authors' contribution:

- A. Study design/planning
- B. Data collection/entry
- C. Data analysis/statistics
- D. Data interpretation
- E. Preparation of manuscript
- F. Literature analysis/search
- G. Funds collection

### Abstract

Gestational diabetes mellitus (GDM) is a pregnancy-related condition characterized by glucose intolerance, significantly impacting both maternal and fetal health. Among the modifiable risk factors, physical inactivity has been identified as a key contributor to GDM. The review synthesizes findings from twelve studies conducted across eight countries, with a combined total of 24,583 participants. The results consistently indicate that women who engage in regular physical activity (PA) before and during early pregnancy have a lower risk of developing GDM. The protective effect is particularly pronounced among those active before conception. Despite the consistent associations, variations in study design, population characteristics, and methods for assessing PA lead to differences in reported outcomes. The findings emphasize the importance of PA as a preventive measure, highlighting its role in prenatal care. However, variability in results across studies points to the need for further research to develop specific guidelines and targeted interventions. Incorporating PA into prenatal programs could enhance maternal and child health by reducing the risk of GDM and its associated long-term metabolic complications.

**Keywords:** physical inactivity, gestational diabetes, preventive health services, pregnancy complications, risk factors

### Introduction

Gestational diabetes mellitus (GDM) is a common metabolic condition identified by glucose intolerance that emerges for the first time during pregnancy [1]. It poses significant health risks for both mothers and their offspring, contributing to adverse pregnancy outcomes and increasing the likelihood of long-term metabolic complications [2]. The etiology and genetic foundations of GDM are not yet fully elucidated, prompting intensive research efforts to identify modifiable risk factors that could inform preventive strategies [3,4].

Among the numerous factors implicated in the pathogenesis of GDM, lifestyle behaviors have garnered considerable attention due to their potential

Tables: 2

Figures: 1

References: 39

Submitted: 2024 Nov 5

Accepted: 2025 Jan 7

Published Online: 2025 Feb 20

Bali S, Singh S, Arora A. Exploring the role of physical inactivity in gestational diabetes mellitus: a systematic review of evidence. *Health Prob Civil.* 2026; 20(2): 186-197. <https://doi.org/10.5114/hpc.2025.146537>

**Address for correspondence:** Seveka Bali, Physiotherapy Department, Punjabi University, NH 64, next to Urban Estate Phase II, 147002 Patiala, Punjab, India, e-mail: sevekaphysio@gmail.com

ORCID: Seveka Bali <https://orcid.org/0000-0002-4447-4844>, Sonia Singh <https://orcid.org/0009-0004-6335-9173>

Copyright: © 2026 Seveka Bali, Sonia Singh, Aashima Arora. This is an Open Access journal, all articles are distributed under the terms of the Creative Commons AttributionNonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited and states its license.

for [5]. Amidst the myriad factors influencing GDM development, physical inactivity, characterized by insufficient regular exercise and sedentary behaviors, emerges as a modifiable lifestyle factor warranting thorough examination [6].

The American College of Obstetrics and Gynecology (ACOG) recommends that pregnant women should engage in at least 150 minutes of moderate-intensity aerobic activity throughout the week, which can be broken down into approximately 30 minutes of exercise on >5 days/week in case of no contraindication [7,8].

Aerobic exercise and resistance training are considered some of the safest and most effective forms of exercise for preventing gastrointestinal motility disorders [9,10].

In contemporary societies, sedentary lifestyles have become increasingly prevalent, driven by technological advancements, urbanization, and changing work patterns [11]. As a result, prevalence of physical inactivity among pregnant women is on the rise, raising concerns about its potential contribution to the escalating rates of GDM worldwide [8,12].

Physical activity (PA) reduces GDM risk through protective effects, enhancing glucose uptake in muscles and decreasing insulin resistance, which is crucial during pregnancy when insulin resistance naturally increases [5]. The effects are primarily mediated by improved insulin sensitivity via GLUT4 translocation, increasing glucose uptake by muscle cells, improving mitochondrial function and fatty acid oxidation, enhances endothelial function [13-16]. Regular exercise reduces inflammatory markers, including C-reactive protein (CRP) and interleukin-6 (IL-6), helping to lower diabetes risk [17,18].

While the beneficial effects of regular PA on metabolic health are well-established in the general population, its specific impact on GDM risk during pregnancy warrants further investigation for devising effective preventive strategies and optimizing maternal and fetal health outcomes [19].

Several observational studies have reported associations between physical inactivity and an increased likelihood of developing GDM. However, the evidence remains heterogeneous, with variations in study designs, population characteristics, and measures of PA complicated interpretation [20].

The multifactorial nature of GDM necessitates a meticulous investigation into the interplay between physical inactivity and other established risk factors, such as age, body mass index, dietary habits, and genetic predispositions. The review uses a systematic approach to delineate the independent contribution of physical inactivity toward GDM onset while discerning potential synergistic or mitigating effects in conjunction with other risk factors.

Ultimately, synthesizing the existing evidence on the association between physical inactivity and GDM will provide clinicians, public health policymakers, and expectant mothers with valuable insights to guide preventive measures and interventions. By elucidating the significance of PA in mitigating the GDM risk, the review endeavors to contribute to the optimization of maternal health and the reduction of adverse pregnancy outcomes.

### **Aim of the work**

The review aims at exploring the role of physical inactivity in the development of GDM by analyzing evidence from diverse study designs.

## Methods

### *Search strategy*

Literature searches were conducted using the databases MEDLINE, Scopus, CINAHL, Embase, and Cochrane Library from March 2009 to March 2024, with an update in August 2024. We used various keyword combinations to account for different search parameters in each database: (“pregnancy” OR “gestation” OR “gestational” OR “maternal” OR “antenatal”) AND (“exercise” OR “activity” OR “PA” OR “physical exercise”) AND (“gestational diabetes” OR “GDM” OR “insulin sensitivity” OR “glucose tolerance” OR “diabetes mellitus” OR “diabetes”). Additionally, bibliographies of relevant articles were examined to ensure a comprehensive collection. Three reviewers independently analyzed and selected potential studies identified through the search strategy.

### *Study selection*

Only human studies published in English were considered. We focused on studies involving pregnant women in their first trimester. Studies were excluded if they involved participants at low risk of GDM, those with pre-existing diabetes, high-risk pregnancies, or if they were not conducted in routine care settings. There were no exclusions based on various international definitions of GDM.


### *Quality assessment*

We used the Mixed Methods Appraisal Tool (MMAT) to assess the quality of the original studies. The MMAT was initially developed in 2006 and revised in 2011 [21]. The tool allows for the concurrent appraisal and description of methodological quality in case-control, cohort, and cross-sectional studies. For our assessment, we used section 3 of the MMAT to evaluate the study quality. Each item is marked as yes, no, and cannot tell. The total number of items rated “yes” is counted to produce an overall score. The score is determined by the number of criteria met divided by four, resulting in a range from 25% (\*) for one criterion met to 100% (\*\*\*\*) for meeting all criteria [21-23].

## Literature review results

### *Identified studies*

A comprehensive search across multiple databases initially identified 120 potentially relevant papers. After screening titles and abstracts and removing duplicates, we retrieved full texts of 50 papers for further review. Following applying various exclusion criteria, we excluded several studies, including one where the author failed to provide additional methodological details. In the end, 12 studies (as shown in Table 2) met the inclusion criteria for the synthesis [20,24-33]. The process of study selection is depicted in Figure 1.



**Figure 1.** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram

### Study characteristics

We individually selected the research works included in the synthesis, considering the eligibility criteria. The included studies explored the relationship between PA and GDM across diverse populations, employing various assessment methods. The studies were conducted in eight countries: the USA, Norway, India, Iran, China, Brazil, Canada, and Spain (Table 1). Most papers were cohort studies, with three being case-control and three being cross-sectional. All the studies included pregnant women, both with and without GDM, comprising a total of 24,583 participants. Most of the research used different questionnaires to assess PA (Table 2), with only two studies utilizing pedometers and armbands in addition to questionnaires [28,34]. Different criteria used for the GDM diagnosis are mentioned in Table 1.

**Table 1.** Study characteristics

Study number	Author(s) and year	Study design	Population	Sample size	PA assessment	Diagnostic criteria	Results
1.	Ramos-Levi et al., 2012 [26]	Cohort study	Pregnant women in Spain	2,194	Questionnaire	Carpenter and Coustan's criteria	Light walking (>30 minutes/day) and sports activities ( $\geq 2$ days/week) reduce the chance of GDM.
2.	Li et al., 2013 [33]	Cross-sectional	Pregnant women in Guangzhou, China	571	IPAQ	IADPSG criteria	13% of the women had GDM, associated with increased intake of local foods and lower PA levels ( $\leq 600$ MET-min / week).
3.	Badon et al., 2017 [20]	Cohort study	Pregnant women in Washington, the USA	3,005	Questionnaire	American Diabetes Association guidelines	PA (>3 METs) reduced the risk of GDM by 21% for each 1-point increase in lifestyle score.
4.	Ramos do Nascimento et al., 2019 [31]	Cross-sectional	Low-income pregnant women in Brazil	544	PPAQ	IADPSG criteria	GDM occurred in 17.4% of participants, among which 61.0 % were physically inactive (<1.5 METs).
5.	Leng et al., 2016 [30]	Cross-sectional	Pregnant women in Tianjin, China	11,450	Questionnaire	IADPSG criteria	Extended periods of sitting were markedly linked to a heightened likelihood of developing GDM.
6.	Amiri et al., 2016 [27]	Case-control	Pregnant women in Babol, Iran	200	PPAQ	Carpenter and Coustan criteria	GDM was more prevalent in women with low total PA, as compared to those with higher PA levels (activities with MET >3 MET).
7.	Chasan-Taber et al., 2010 [25]	Cohort study	Hispanic women in Massachusetts, the USA	632	PPAQ	American Diabetes Association criteria	18.7% of participants met the guidelines for PA before pregnancy, while 5.2% met these guidelines in early pregnancy.
8.	Mørkrid et al., 2014 [34]	Cohort study	Pregnant women in Oslo, Norway	759	Sense Wear™ Armband	IADPSG criteria	30% of women with GDM reported engaging in regular PA before pregnancy, as compared to 44% of those without GDM.

Study number	Author(s) and year	Study design	Population	Sample size	PA assessment	Diagnostic criteria	Results
9.	Mishra et al., 2020 [32]	Case-control	Pregnant women in Karnataka, India	373	IPAQ	Carpenter and Coustan criteria, later switched to DIPSI criteria	Total PA <600 MET-minutes/week is a significant risk factor for GDM.
10.	Retnakaran et al., 2009 [24]	Observational cohort study	Healthy pregnant women in Canada	851	Baecke questionnaire	GCT and OGTT	Women with glucose tolerance were engaged in higher levels of vigorous/sports activity before pregnancy than those with abnormal glucose tolerance.
11.	Anjana et al., 2016 [28]	Case-control	Pregnant women in Chennai, India (GDM cases and controls)	795	MPAQ, pedometer	IADPSG criteria	Sedentary behavior increased the likelihood of GDM four times, whereas walking reduced GDM by 70%.
12.	Badon et al., 2016 [29]	Cohort study	Pregnant women in Washington State, the United States	3,209	LTPA	Carpenter and Coustan's criteria	High LTPA duration before pregnancy (6-32.5 hrs/wk), lowers the risk of GDM by 45%.

Notes: IADPSG – The International Association of Diabetes and Pregnancy Study Groups, GCT – Glucose Challenge Test, OGTT – Oral Glucose Tolerance Test, DIPSI – Diabetes in Pregnancy Study Group India, MPAQ – Madras Diabetes Research Foundation – Physical Activity Questionnaire, IPAQ – International Physical Activity Questionnaire, PPAQ – Pregnancy Physical Activity Questionnaire, LTPA – Leisure-Time Physical Activity.

### Methodological quality

The studies' quality was assessed using the MMAT, which recommended one moderate-quality study and eleven high-quality studies (Table 2). Any disagreement was resolved by collaborative deliberation.

**Table 2.** Quality assessment

Study number	Study	Representative participants	Appropriate measurements	Confounding factors controlled	Complete follow-up	Exposures administered as intended	Overall score	Quality of the study
1.	Badon et al., 2016 [29]	Yes	Yes	Yes	Not mentioned	Yes	80%	Moderate to High-Quality Study
2.	Retnakaran et al., 2009 [24]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
3.	Anjana et al., 2016 [28]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
4.	Ramos-Levi et al., 2012 [26]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study

Study number	Study	Representative participants	Appropriate measurements	Confounding factors controlled	Complete follow-up	Exposures administered as intended	Overall score	Quality of the study
5.	Badon et al., 2017 [20]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
6.	Ramos do Nascimento et al., 2019 [31]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
7.	Leng et al., 2016 [30]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
8.	Amiri et al., 2016 [27]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
9.	Chasan-Taber et al., 2010 [25]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
10.	Mørkrid et al., 2014 [34]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
11.	Mishra et al., 2020 [32]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study
12.	Li et al., 2013 [33]	Yes	Yes	Yes	Yes	Yes	100%	High-Quality Study

### *PA during pregnancy*

To date, three observational studies have analyzed the relationship between pre-pregnancy PA levels and glucose and insulin metabolism during pregnancy. Pregnant women from Norway were studied to find the link between pre-pregnancy PA measured objectively and GDM. Interestingly, only 30% of women with GDM reported engaging in regular PA before pregnancy, as compared to 44% of those without GDM [34]. One research focused on PA types and metabolic parameters during pregnancy. Glucose intolerance decreased with a higher level of pre-pregnancy total PA ( $p=0.024$ ). Women with standard glucose tolerance reported significantly higher levels of vigorous/sports activity before pregnancy than those with abnormal glucose tolerance ( $p=0.004$ ) [24]. Additionally, a prospective cohort study among Hispanic women in Massachusetts evaluated PA, psychosocial stress, and GDM risk, finding that 18.7% of participants met the guidelines for PA before pregnancy, while 5.2% met the guidelines in early pregnancy (30 minutes of moderate PA on >5days/week) [25].

Two studies have objectively measured PA levels to examine their association with glucose intolerance in pregnancy. One case-control study involving 795 pregnant women used pedometers and a questionnaire to assess PA status. The results demonstrate that women with GDM exhibit significantly higher levels of sedentary behavior, as compared to women without the condition (86.2% vs. 61.2%;  $p<0.001$ ). Sedentary behavior demonstrated a significant fourfold increase in the likelihood of adverse outcomes, whereas engaging in recreational walking was linked to a notable 70% reduction in the risk of GDM [28]. The study conducted by Morkrid et al. [34] utilized armbands to monitor PA during early gestation. Women diagnosed with GDM were found to take fewer steps on average per day (Mean 7964 steps/day) and engage in moderate intensity of PA (3-6 METs;  $p<0.005$ ), as compared to women without GDM [34].

Three observational researches have evaluated the effect of PA during pregnancy on GDM development. Authors used the questionnaire to assess PA levels in early pregnancy and found that GDM was more

prevalent in women with low total PA ( $p=0.001$ ), as compared to those reporting higher PA levels (activities with MET  $>3$  MET) [27]. A study in China examined the relationship between dietary habits, PA, and cognitive factors with GDM risk, finding that 13% of the women had GDM, associated with increased intake of local foods and lower PA levels ( $\leq 600$  MET-min/week) [33]. In Karnataka, India, pooled data identified physical inactivity (total PA  $<600$  MET-min/week) as a significant risk factor for GDM (adjusted OR: 21.0;  $p=0.003$ ) [32].

A cross-sectional and cohort study involving 11,450 pregnant women in China and 3,005 women in Washington supported the findings. Women with GDM were more likely to be engaged in less than three hours of LTPA and less than one hour of household work. Pregnant women diagnosed with GDM demonstrated a reduced propensity to participate in moderate to vigorous PA during their pregnancy (79.8% vs. 81.6%;  $p=0.191$ ) [30]. In Washington, adherence to a healthy lifestyle, including PA ( $>3$  METs) reduced the risk of GDM by 21% for each 1-point increase in lifestyle score [20]. Additionally, a cohort study from Spain found that light walking ( $>30$  min/day) and sports activities ( $\geq 2$  days/week) reduce the chances of GDM [26]. One study investigated LTPA and its association with GDM risk, finding that if LTPA duration before pregnancy is high (6-32.5 hrs/wk), it lowers the risk of GDM by 45% [29].

In Brazil, a study of 544 low-income pregnant women found that GDM occurred in 17.4% of participants, among which 61.0% were physically inactive ( $<1.5$  METs) [31]. Cumulative results suggest that engaging in PA before and during the early stages of pregnancy is correlated with a decreased likelihood of GDM development, with the most robust protective impact noted in women who were physically active before conception.

#### *Discussion of the review results*

The findings of the systematic review provide a comprehensive assessment of the PA as a risk factor for GDM. The evidence underscores the potential benefits of PA as a preventive measure for GDM but also highlights the complexity and variability in outcomes across different studies. The effectiveness of PA programs in controlling GDM was mentioned in a systematic review [35], whereas another systematic review and meta-analysis revealed a curved pattern of influence between activity done before pregnancy and the risk of GDM [9]. Our findings align with the broader literature highlighting the beneficial effects of PA on overall metabolic health [6]. The relationship between PA and GDM is crucial, as highlighted by those who discussed the association between physical inactivity and mortality due to diabetes mellitus [36]. Furthermore, physical inactivity is also a leading risk factor for various health conditions, including GDM [37]. A meta-analysis revealed a significant association between elevated PA levels before pregnancy and a reduced likelihood of developing GDM [9]. Activities performed before pregnancy seem to have a more pronounced protective effect, as compared to those initiated during pregnancy. It may be due to the establishment of better metabolic health before pregnancy [38]. The importance of lifestyle modifications, including PA, in improving outcomes for women with GDM was underscored in another study [39]. Increased sedentary time, regardless of PA levels, is also associated with a higher risk of GDM. It suggests that reducing sedentary time is as important as promoting PA [30]. Low income was identified as a contributing factor to physical inactivity among females, which subsequently increases the risk of developing GDM [31]. In conclusion, physical inactivity plays a significant role in the development and management of GDM. Incorporating PA interventions in the care of pregnant women can potentially reduce the risk of GDM and improve maternal and child health outcomes.

Despite the overall positive association, the review highlighted significant heterogeneity in the outcomes of the studies. Self-reported PA data could be susceptible to recall bias. Future research should consider using objective measures of PA, such as accelerometers, to validate the findings. Standardizing the measurement of PA and sedentary behavior will improve comparability across studies. Variations in study designs, population characteristics, and PA assessment methods contribute to inconsistent findings. For example, while some studies reported a substantial risk reduction, others found minimal effect. Additionally, longitudinal studies tracking PA patterns from preconception through postpartum would provide valuable insights into the timing and duration of activity needed to confer protective effects.

## Conclusions

PA holds promise as a modifiable risk factor for GDM, with evidence supporting its role in prevention. However, variability in findings necessitates further research to refine recommendations and develop effective intervention strategies tailored to diverse populations.

## Disclosures and acknowledgments

The authors declare no conflicts of interest with respect to the research, authorship, and/or publication of this article.

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Artificial intelligence (AI) was not used in the creation of the manuscript.

## References:

1. Moon JH, Jang HC. Gestational diabetes mellitus: diagnostic approaches and maternal-offspring complications. *Diabetes Metab J.* 2022; 46: 3-14. <https://doi.org/10.4093/dmj.2021.0335>
2. Damm P, Houshmand-Oeregaard A, Kelstrup L, Lauenborg J, Mathiesen ER, Clausen TD. Gestational diabetes mellitus and long-term consequences for mother and offspring: a view from Denmark. *Diabetologia.* 2016; 59: 1396-99. <https://doi.org/10.1007/s00125-016-3985-5>
3. Chen L, Mayo R, Chatry A, Hu G. Gestational diabetes mellitus: its epidemiology and implication beyond pregnancy. *Current Epidemiology Reports.* 2016; 3: 1-11. <https://doi.org/10.1007/s40471-016-0063-y>
4. Ray GW, Zeng Q, Kusi P, Zhang H, Shao T, Yang T, et al. Genetic and inflammatory factors underlying gestational diabetes mellitus: a review. *Frontiers in Endocrinology.* 2024; 15: 1399694. <https://doi.org/10.3389/fendo.2024.1399694>
5. Rasmussen L, Poulsen CW, Kampmann U, Smedegaard SB, Ovesen PG, Fuglsang J. Diet and healthy lifestyle in the management of gestational diabetes mellitus. *Nutrients.* 2020; 12: 1-24. <https://doi.org/10.3390/nu12103050>
6. Zhang C, Ning Y. Effect of dietary and lifestyle factors on the risk of gestational diabetes: review of epidemiologic evidence. *Am J Clin Nutr.* 2011; 94(6): S1975-9. <https://doi.org/10.3945/ajcn.110.001032>
7. Belachew DZ, Melese T, Negese K, Abebe GF, Kassa ZY. Antenatal physical exercise level and its associated factors among pregnant women in Hawassa City, Sidama Region, Ethiopia. *PLoS One.* 2023; 18(4): e0280220. <https://doi.org/10.1371/journal.pone.0280220>
8. Leiferman JA, Gutilla MJ, Paulson JF, Pivarnik J. Antenatal physical activity counseling among healthcare providers. *Open J Obstet Gynecol.* 2012; 02(04): 346-55. <https://doi.org/10.4236/ojog.2012.24073>

9. Aune D, Sen A, Henriksen T, Saugstad OD, Tonstad S. Physical activity and the risk of gestational diabetes mellitus: a systematic review and dose-response meta-analysis of epidemiological studies. *Eur J Epidemiol.* 2016; 31(10): 967-97. <https://doi.org/10.1007/s10654-016-0176-0>
10. Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *The Lancet.* 2012; 380(9838): 219-229. [https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)
11. Kintiraki E, Mintziori G, Goulis D.G. Pathogenesis of gestational diabetes mellitus. In: Rodriguez-Saldana J, editor. *The diabetes textbook.* Cham: Springer; 2023. p. 247-259. [https://doi.org/10.1007/978-3-031-25519-9\\_15](https://doi.org/10.1007/978-3-031-25519-9_15)
12. Harreiter J, Dovjak G, Kautzky-Willer A. Gestational diabetes mellitus and cardiovascular risk after pregnancy. *Women's Health.* 2014; 10: 91-108. <https://doi.org/10.2217/WHE.13.69>
13. Jung S-R, Lee J-H, Ryu H, Gao Y, Lee J. Lithium and exercise ameliorate insulin-deficient hyperglycemia by independently attenuating pancreatic  $\alpha$ -cell mass and hepatic gluconeogenesis. *Korean Journal of Physiology and Pharmacology.* 2024; 28(1): 31-38. <https://doi.org/10.4196/kjpp.2024.28.1.31>
14. Bowman PRT, Smith GL, Gould GW. Run for your life: can exercise be used to effectively target GLUT4 in diabetic cardiac disease?. *PeerJ.* 2021; 9: e11485. <https://doi.org/10.7717/peerj.11485>
15. Langlois A, Forterre A V, Pinget M, Bouzakri K. Impact of moderate exercise on fatty acid oxidation in pancreatic B-Cells and skeletal muscle. *J Endocrinol Invest.* 2021; 44(9): 1815-1825. <https://doi.org/10.1007/s40618-021-01551-2>
16. Hjerkind KV, Stenehjem JS, Nilsen TIL. Adiposity, physical activity and risk of diabetes mellitus: prospective data from the population-based HUNT study, Norway. *BMJ Open.* 2017; 7(1): e013142. <https://doi.org/10.1136/bmjopen-2016-013142>
17. Mutie PM, Drake I, Ericson U, Teleka S, Schulz CA, Stocks T, et al. Different domains of self-reported physical activity and risk of type 2 diabetes in a population-based Swedish cohort: the Malmö diet and cancer study. *BMC Public Health.* 2020; 20(1). <https://doi.org/10.1186/s12889-020-8344-2>
18. Abdelrahim HM, Madani AM, El-Hadiyah TM, Mahmoud AN. Exercise as component of standard diabetes management, the challenges to improve performance among sudanese type 2 diabetes. *Open J Endocr Metab Dis.* 2020; 10(08): 107-116. <https://doi.org/10.4236/ojemd.2020.108011>
19. Koivusalo SB, Rönö K, Klemetti MM, Roine RP, Lindström J, Erkkola M, et al. gestational diabetes mellitus can be prevented by lifestyle intervention: the Finnish Gestational Diabetes Prevention Study (RADIEL): a randomized controlled trial. *Diabetes Care.* 2016; 39(1): 24-30. <https://doi.org/10.2337/dc15-0511>
20. Badon SE, Enquobahrie DA, Wartko PD, Miller RS, Qiu C, Gelaye B, et al. Healthy lifestyle during early pregnancy and risk of gestational diabetes mellitus. *Am J Epidemiol.* 2017; 186: 326-333. <https://doi.org/10.1093/aje/kwx095>
21. Pace R, Pluye P, Bartlett G, Macaulay AC, Salsberg J, Jagosh J, et al. Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool (MMAT) for systematic mixed studies review. *Int J Nurs Stud.* 2012; 49(1): 47-53. <https://doi.org/10.1016/j.ijnurstu.2011.07.002>
22. Hong QN, Fàbregues S, Bartlett G, Boardman F, Cargo M, Dagenais P, et al. The Mixed Methods Appraisal Tool (MMAT) version 2018 for information professionals and researchers. *Education for Information.* 2018; 34(4): 285-291. <https://doi.org/10.3233/EFI-180221>

23. Peterson J, Welch V, Losos M, Tugwell PJ. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa: Ottawa Hospital Research Institute; 2011.
24. Retnakaran R, Qi Y, Sermer M, Connelly PW, Zinman B, Hanley AJG. Pre-gravid physical activity and reduced risk of glucose intolerance in pregnancy: the role of insulin sensitivity. *Clin Endocrinol (Oxf)*. 2009; 70: 615-22. <https://doi.org/10.1111/j.1365-2265.2008.03393.x>
25. Chasan-Taber L, Fortner RT, Gollenberg A, Buonnaccorsi J, Dole N, Markenson G. A prospective cohort study of modifiable risk factors for gestational diabetes among Hispanic women: design and baseline characteristics. *Journal of Women's Health*. 2010; 19: 117-124. <https://doi.org/10.1089/jwh.2009.1416>
26. Ramos-Leví AM, Pérez-Ferre N, Fernández MD, Del Valle L, Bordiu E, Bedia AR, et al. Risk factors for gestational diabetes mellitus in a large population of women living in Spain: implications for preventative strategies. *Int J Endocrinol*. 2012; 2012: 312529. <https://doi.org/10.1155/2012/312529>
27. Nasiri-Amiri F, Bakhtiari A, Faramarzi M, Rad HA, Pasha H. The association between physical activity during pregnancy and gestational diabetes mellitus: a case-control study. *Int J Endocrinol Metab*. 2016; 14: 37123. <https://doi.org/10.5812/ijem.37123>
28. Anjana RM, Sudha V, Lakshmipriya N, Anitha C, Unnikrishnan R, Bhavadharini B, et al. Physical activity patterns and gestational diabetes outcomes – the wings project. *Diabetes Res Clin Pract*. 2016; 116: 253-262. <https://doi.org/10.1016/j.diabres.2016.04.041>
29. Badon SE, Wartko PD, Qiu C, Sorensen TK, Williams MA, Enquobahrie DA. Leisure time physical activity and gestational diabetes mellitus in the Omega Study. *Med Sci Sports Exerc*. 2016; 48(6): 1044. <https://doi.org/10.1249/MSS.0000000000000866>
30. Leng J, Liu G, Zhang C, Xin S, Chen F, Li B, et al. Physical activity, sedentary behaviors and risk of gestational diabetes mellitus: a population-based cross-sectional study in Tianjin, China. *Eur J Endocrinol*. 2016; 174: 763-773. <https://doi.org/10.1530/EJE-15-1103>
31. Ramos do Nascimento G, do Carmo Borges M, Figueiroa JN, Alves LV, Alves JG. Physical activity pattern in early pregnancy and gestational diabetes mellitus risk among low-income women: a prospective cross-sectional study. *SAGE Open Med*. 2019; 7. <https://doi.org/10.1177/2050312119875922>
32. Mishra S, Shetty A, Rao CR, Nayak S, Kamath A. Risk factors for gestational diabetes mellitus: a prospective case-control study from coastal Karnataka. *Clin Epidemiol Glob Health*. 2020; 8(4): 1082-1088. <https://doi.org/10.1016/j.cegh.2020.03.024>
33. Li Q, Xiong R, Wang L, Cui J, Shi L, Liu Y, et al. Associations of dietary habits, physical activity and cognitive views with gestational diabetes mellitus among Chinese women. *Public Health Nutr*. 2014; 17: 1850-1857. <https://doi.org/10.1017/S1368980013001882>
34. Mørkrid K, Jenum AK, Berntsen S, Sletner L, Richardsen KR, Vangen S, et al. Objectively recorded physical activity and the association with gestational diabetes. *Scand J Med Sci Sports*. 2014; 24: e389-97. <https://doi.org/10.1111/sms.12183>
35. Laredo-Aguilera JA, Gallardo-Bravo M, Rabanales-Sotos JA, Cobo-Cuenca AI, Carmona-Torres JM. Physical activity programs during pregnancy are effective for the control of gestational diabetes mellitus. *International Journal of Environmental Research and Public Health*. 2020; 17: 6151. <https://doi.org/10.3390/ijerph17176151>
36. Silva DAS, Naghavi M, Duncan BB, Schmidt MI, De Souza MDFM, Malta DC. Physical inactivity as risk factor for mortality by diabetes mellitus in Brazil in 1990, 2006, and 2016. *Diabetol Metab Syndr*. 2019; 11: 1-11. <https://doi.org/10.1186/s13098-019-0419-9>

37. Hartsock E, Huynh A. Lifestyle modifications may improve glycemic control in pregnant women. *J Comp Nurs Res Care*. 2020; 5: JCNRC-163. <https://doi.org/10.33790/jcnrc1100163>
38. Tobias DK, Zhang C, Van Dam RM, Bowers K, Hu FB. Physical activity before and during pregnancy and risk of gestational diabetes mellitus: a meta-analysis. *Diabetes Care*. 2011; 34: 223. <https://doi.org/10.2337/dc10-1368>
39. Marchetti D, Carrozzino D, Fraticelli F, Fulcheri M, Vitacolonna E. Quality of life in women with gestational diabetes mellitus: a systematic review. *J Diabetes Res*. 2017; 2017: 7058082. <https://doi.org/10.1155/2017/7058082>